

Early-type Galaxies in the Hubble Deep Field.

The $\langle \mu_e \rangle - r_e$ relation and the lack of large galaxies at high redshift

Giovanni Fasano

Osservatorio Astronomico di Padova, vicolo dell'Osservatorio 5, 35122 Padova, Italy

Stefano Cristiani

Dipartimento di Astronomia, Università di Padova, vicolo dell'Osservatorio 5, 35122 Padova, Italy

Stephane Arnouts

Osservatorio Astronomico di Padova, vicolo dell'Osservatorio 5, 35122 Padova, Italy

Institut d'Astrophysique de Paris, 98-bis Boulevard Arago, 75014 Paris

and

Michele Filippi

Dipartimento di Astronomia, Università di Padova, vicolo dell'Osservatorio 5, 35122 Padova, Italy

Received _____; accepted _____

ABSTRACT

We present the results of the detailed surface photometry of a sample of early-type galaxies in the Hubble Deep Field. Effective radii, surface brightnesses and total V_{606} magnitudes have been obtained, as well as U_{300} , B_{450} , I_{814} , J , H and K colors, which are compared with the predictions of chemical-spectrophotometric models of population synthesis. Spectroscopic redshifts are available for 23 objects. For other 25 photometric redshifts are given. In the $< \mu_e > -r_e$ plane the early-type galaxies of the HDF, once the appropriate $K + E$ corrections are applied, turn out to follow the ‘*rest frame*’ Kormendy relation. This evidence, linked to the dynamical information gathered by Steidel et al. (1996), indicates that these galaxies, even at $z \simeq 2 - 3$, lie in the Fundamental Plane, in a virial equilibrium condition. At the same redshifts a statistically significant lack of large galaxies (i.e. with $\log r_e^{kpc} > 0.2$) is observed.

Subject headings: cosmology: observations - galaxies: photometry - galaxies: evolution
- galaxies: elliptical and lenticular

1. Introduction

The formation and evolution of galaxies is a complex process, which may involve starbursts, infall, interactions (e.g., Dressler et al. 1994a,b; Lilly et al. 1995; Moore et al. 1996). The Tully-Fisher (1977) and the Fundamental Plane (FP) (Dressler et al. 1987a; Djorgovski & Davis 1987) relations are valuable sources of information for disk and elliptical galaxies, respectively. In particular, the FP and its projection on the average surface brightness ($\langle \mu_e \rangle$) effective radius (r_e) plane (Kormendy relation, Kormendy 1985) have been used as powerful tools for estimating distances of galaxy clusters (Dressler et al. 1987b, Lynden-Bell et al. 1988, Faber et al. 1989) and, more recently, for investigating the evolution of elliptical galaxies (Bender et al. 1992, Pahre and Djorgovski 1996, Fasano et al. 1997).

If the luminosity profiles and the dynamical structures of the galaxies are similar, then the very existence of the FP implies a virial equilibrium condition. Moreover, the slope of the FP (which slightly differs from that expected in the case of ‘pure’ virial equilibrium) could imply that the M/L ratio depends on the mass of the galaxy (Djorgovski and Santiago 1993). Recent advances in the sensitivity and resolution of the observations both in imaging and spectroscopy with the *Hubble Space Telescope* (HST) and from the ground have greatly enlarged the horizon of morphological and dynamical studies, thus allowing the investigation of the FP in clusters with redshift up to $z \sim 0.6$ (Kelson et al. 1997). The backward evolution of the surface brightness derived from the zero point of the FP in these clusters seems consistent with a passive evolution of galaxies (Bender et al. 1997). On the other hand, it is still matter of debate whether the slope of the FP (i.e. the function $M/L = f(M)$) varies with the redshift (van Dokkum and Franx 1996, Kelson et al. 1997, Jørgensen and Hjorth 1997). The difficulty of obtaining reliable velocity dispersion estimates represents the key obstacle in trying to improve the results of FP studies, as well as to extend them to higher redshifts. In spite of its larger scatter, the Kormendy relation $\langle \mu_e \rangle - r_e$ allows to overcome this limitation (Kjærgaard et al. 1993, Pahre et al. 1996, Fasano et al. 1997). Actually, from deep HST imaging of fields around powerful radio-sources and radio-loud quasars, Dickinson (1995, 1997a) was able to obtain the Kormendy relation of clusters

up to $z \sim 1.2$.

The Hubble Deep Field (Williams et al. 1996, see also Ellis 1997; hereafter HDF) is perhaps the most impressive example of the recent advances in optical imaging. The very long exposure time, the observational procedures and the data handling techniques of HDF have provided exceptional spatial resolution and depth. The unprecedented low surface brightness level reached by the HDF data turns out to be particularly suited for performing detailed surface photometry of objects whose brightness slowly decreases towards the outer regions, as in the case of elliptical galaxies.

We have undertaken a long-term project aimed at producing detailed surface photometry in the four available bands of a sample of early-type galaxies in the HDF. The main scientific goal of this project is to put strong observational constraints to the theories modeling the evolution of elliptical galaxies (e.g. Tantalo et al. 1996, Kauffmann and Charlot 1997).

The detailed surface photometry of our galaxy sample in the V_{606} band is presented elsewhere (Fasano and Filippi 1998; hereafter FF98). In the present paper we discuss the improvements that the morphological information, together with the photometry in different optical and infrared bands, can produce in understanding the processes of galaxy formation and evolution.

In Section 2 we report on the database we use to perform the present analysis, referring to FF98 as far as the original sample selection and the detailed surface photometry are concerned. In the same section we describe of the auxiliary data (taken from both network and recent literature) useful for the present analysis. In Section 3 the technique we used to extract from the luminosity profiles the relevant photometric parameters is discussed. In Section 4 we analyze the relation $\langle \mu_e \rangle - r_e$ (Kormendy relation; hereafter KR) for early-type galaxies in the *HDF*. In the same section we discuss the K +evolutionary corrections expected from the current models on the basis of the available optical and infrared photometry, and derive photometric redshifts for the galaxies without spectroscopic information. Discussion and conclusions are given in Section 5. Unless otherwise noted we assume $H_o = 50$ km/s/Mpc and $q_o = 0.5$ throughout.

2. The database and the sample selection

The present sample of *early-type* galaxies has been extracted from the second release of the *WFPC2 – HDF* frames, in the V_{606} band. The selection is based on the photometry carried out by Couch (1996) with the automated SExtractor algorithm (Bertin and Arnouts 1996).

A preliminary list of candidates was defined including all the HDF objects satisfying the following criteria: *a)* Kron AB magnitude in the V_{606} band ≤ 26.3 ; *b)* number of pixels above the threshold limit of 1.3σ of the background noise ≥ 200 ; *c)* star/galaxy classifier (s/g) ≤ 0.6 ($s/g=1$ means 'star'). The combination of the first two limits ensures a sufficient signal to noise ratio and an adequate number of independent isophotes to carry out a reliable morphological analysis.

Each of the 401 objects matching the above limits was examined with the *IMEXAM-IRAF* tool to produce a first screening against late-type objects. The preliminary morphological classification was also compared with the ones by Van den Bergh et al. (1996) and Statler (1996), finding a general good agreement. Detailed surface photometry (luminosity and geometrical profiles) was carried out on the resulting list of 162 *early-type* candidates. From the analysis of the luminosity and geometrical profiles, 34 objects of the sample have been recognized to be '*disk-dominated*' objects (likely *Sa* galaxies), whereas 28 more objects showed peculiar or unclassifiable profiles. These galaxies were excluded from the final sample of 99 '*bona fide*' *early-type* galaxies.

Further details about the sample selection and surface photometry are given in FF98. We recall here that, on the basis of their luminosity profiles, our *HDF early-type* galaxies were divided in three different classes: (FF98; see also Fasano et al. 1996): *1)* the '*Normal*' class (54 galaxies), for which the de Vaucouleurs law is well sampled up to the innermost isophote that is not significantly affected by the *PSF*; *2)* the '*Flat*' class (36 galaxies), characterized by an inward flattening of the luminosity profiles (with respect to the de Vaucouleurs' law) which cannot be ascribed to the effect of the *PSF*; *3)* the '*Merger*' class (9 galaxies), in which isophotal contours show the existence of complex inner structures (two or more nuclei) embedded inside common

envelopes, roughly obeying the $r^{1/4}$ law.

It is worth to point out that the morphological classification of galaxies in our sample is necessarily much more a qualitative than a quantitative one. In most cases the spatial resolution is ‘critical’ and many faint galaxies appear heavily disturbed by the presence of close (often multiple) companions or peculiar structures. Sometimes the only unquestionable characteristic of the object is the presence of a strong spheroidal component. For this reason, although the main requirement for final inclusion in our sample is the prominence of the $r^{1/4}$ component in the luminosity profile, we consider more appropriate referring to our galaxies as ‘*early-type*’.

An additional problem concerning the morphological classification comes from the fact that the light distribution of very high redshift galaxies in the V_{606} band (rest-frame mid and far-IV wavelengths) may reflect the comparatively chaotic distribution of star-formation rather than the structural morphology of galaxies. In Section 5 this point will be addressed in more detail.

In the following we will restrict our analysis to the galaxies for which a spectroscopic redshift is available (23), or a reliable redshift estimate is possible (25), on the basis of multi-band (optical and infrared) photometry (see Section 4.2).

In Table 1 we report the relevant data for this sub-sample: in column (1) the identification given by Williams et al. (1996); in columns (2) and (3) the coordinates; in column (4) the luminosity profile class; in columns (5) and (6) the effective radius and surface brightness derived from the V_{606} photometry.

In column (9) the total V_{606} magnitude, estimated with the detailed surface photometry of FF98, is reported. In order to obtain the total magnitudes in the other three WFPC2 bands we have followed an indirect procedure. We have obtained SExtractor magnitudes in all four bands using for each galaxy of the sample an optimized aperture derived from the V_{606} image. Then the SExtractor magnitudes have been scaled by the offset between the total and the SExtractor V_{606} magnitudes. This procedure has been adopted because for galaxies with extended profiles like ellipticals the total magnitude derived from the detailed surface photometry (V_{606} band) is

more reliable than the ones provided by any automated photometry. As an example, Fig. 1 shows the biases of the SExtractor total magnitudes as a function of the galaxy size and surface brightness. The color photometry scheme we used is a good approximation if the galaxies do not have significant color gradients. This is true for '*Normal*', nearby ellipticals, where the color gradients turn out to be usually lower than few tenths of magnitude (Franx and Illingworth 1990, Peletier et al. 1990). In Section 5 we will see that also star-forming galaxies at intermediate redshift seem not to show significant color gradients. In any case, the errors expected as due to possible color gradients should be quite lower than the systematic uncertainties associated with automated magnitude estimates (see fig.1). The total U_{300} , B_{450} and I_{814} magnitudes are reported in Table 1 (columns 7, 8 and 10, respectively).

Dickinson (1997b) has provided deep, near infrared (J, H, K) images of the *HDF* taken with the *KPNO – IRIM* camera at the Mayall 4-m telescope (total exposure time of ~ 45 hours). The total J , H and K magnitudes, again estimated with the SExtractor algorithm, are reported in columns (11), (12) and (13), respectively. Column (14) is a flag defining the reliability of the infrared photometry.

The redshifts of the galaxies are reported in column (15). The photometric redshifts are in parentheses, and their uncertainties are given in column (16).

3. Deriving r_e and $\langle \mu_e \rangle$

Many galaxies in the present sample show a luminosity profile close to the instrumental PSF. A restoration procedure is needed to extract useful morphological information. Fasano et al.(1997) have shown that the '*Multi-Gaussian Expansion*' deconvolution technique (*EMGDEC*, Bendinelli 1991) gives good results in recovering the '*true*' equivalent half-light radius r_e and the corresponding average surface brightness $\langle \mu_e \rangle$ from ground-based observations of elliptical galaxies in distant clusters. In particular, by using numerical simulations of synthetic elliptical galaxies, Fasano et al.(1997) concluded that this deconvolution technique is efficient down to

$r_e^{true} \simeq FWHM$. The input data of the *EMGDEC* algorithm are the ‘*equivalent*’ luminosity profiles of the galaxy and of the *PSF*, both represented in analytical form by means of suitable series of gaussians.

In the case of the present sample of *HDF* galaxies, there are particular problems inherent to the data: (a) the *PSF* profile is known not to be a decreasing monotonic function of radius. This implies that, in order to give a good representation of the *PSF*, we should use a multi-gaussian expansion with free gaussian centers (i.e. increased number of parameters); (b) the *PSF* profile exhibits quite extended wings which can in principle affect the total magnitude estimate if some analytic law is assumed to extrapolate the luminosity profiles; (c) the presence of strong diffraction spikes in the *PSF* can alter the shape of the isophotes, thus affecting the profiles especially in case of very bright galactic nuclei; (d) the shape of the *PSF* is well known to depend on the position inside the different *WFPC* frames.

Concerning the point (a), we checked both the ‘*free-center*’ and the ‘*fixed-center*’ multi-gaussian representation of the *PSF* for a few typical cases. We found negligible differences in the deconvolved profiles and no differences in the effective radii between the two procedures. Since the ‘*free-center*’ approach turns out to be much more time consuming with respect to the other one, we decided to ignore the characteristic oscillations of the *PSF* profile in the multi-gaussian representation of the *PSF*.

The points (c) and (d) can be important when dealing with two-dimensional problems (modeling and removal of point sources, Lucy deconvolution, etc.). They can be neglected in our case, since the *EMGDEC* deconvolution operates on ‘*equivalent*’ (radially averaged) profiles.

The point (b) is a delicate one, since it becomes more and more important as the galaxy size becomes comparable with the *PSF* size. In this case it is necessary to impose that the multi-gaussian extrapolation of the outer galaxy profile does not fall below the *PSF* profile itself. To this end we forced the extrapolation of the luminosity profiles of very small galaxies to converge smoothly towards the *PSF* profile at large radii. Luminosity profiles with an effective radius larger than three times the *FWHM* were extrapolated simply with a de Vaucouleurs’ law

(see FF98).

Although preserving the total luminosity, the *EMGDEC* deconvolution is not univocal. Actually, it depends on the multi-gaussian representation (and extrapolation) of the profiles, as well as on the so called ‘*regularization*’ technique (see Bendinelli 1991 for details). However, we found that changing these assumptions within wide ranges produces only marginal changes in the resulting deconvolved profiles and no significant changes at all in the corresponding half-light (*effective*) radii r_e . The deconvolution turns out to be quite robust as far as the estimation of the equivalent effective radius r_e is concerned. It is also worth stressing that the morphological classification (and then the inclusion in the sample) is decided before deconvolution. In other words, the profile shape uncertainties associated with the *EMGDEC* technique do not affect the sample selection.

Fig. 2 shows the application of this algorithm in four extreme cases. The comparison between the effective radii before and after deconvolution is illustrated in Figure 3.

In order to explore how reliable is the *EMGDEC* technique in our particular case, we produced a set of simulations specifically designed to reproduce the observational conditions of our sample of *HDF* ellipticals (*FWHM*/pixel-size ratio, noise, range of r_e /*FWHM* ratios, range of magnitudes). Several frames of synthetic elliptical galaxies, with different magnitudes and effective radii, were first convolved with the theoretical *PSF* of the *WFPC2*, and then were processed using our techniques for the extraction and deconvolution of luminosity profiles. Figure 3 illustrates the results of this analysis. In particular, the bias in the r_e measured in the simulations (solid line) matches the distribution of the observed *HDF* ellipticals (open circles). Figure 3 confirms the previous finding by Fasano et al.(1997) that the ‘*true*’ effective radius can be confidently recovered down to r_e /*FWHM* $\simeq 1$ (see filled circles).

In order to obtain reliable error estimates for r_e , we have also to take into account the uncertainties in the total V_{606} magnitudes. Adopting the errors provided in this band by the automated SExtractor photometry ($\sim 0^m.1$ on average; $\sim 0^m.35$ in the worst cases), as a conservative estimate for our detailed surface photometry, we can estimate the log-error in r_e to

be ~ 0.02 on average (~ 0.06 in the worst cases). The radius r_e and the average *effective* surface brightness $\langle \mu_e \rangle$, listed in columns 5 and 6 of Table 1, are values estimated in the V_{606} band after deconvolution.

4. The Kormendy relation of *HDF* ellipticals

The relation between the effective radius r_e and the corresponding average surface brightness $\langle \mu_e \rangle$ (Kormendy 1985) is just a projection of the Fundamental Plane of elliptical galaxies (Djorgovski & Davis 1987).

The zero point of $\langle \mu_e \rangle$ in the Kormendy relation (hereafter *KR*) has been recently used in galaxy clusters at different redshifts, as a standard candle to perform the classical Tolman test for the expansion of the Universe and for testing the evolutionary models of galaxies (Pahre et al. 1996, Fasano et al. 1997).

In the deep fields (in particular in the *HDF*) we are dealing with a completely different situation: most of the galaxies are not in clusters and have very different redshifts. Still, putting the galaxies in the $(\langle \mu_e \rangle - r_e)$ plane is an interesting exercise, as we will show in the following.

Figure 4 shows the $\langle \mu_e \rangle - r_e$ plane for the 23 *HDF* early-type galaxies in our sample with spectroscopic redshift. The effective radii (in kpc) are computed from the deconvolved luminosity profiles. The triangles in the plots represent the galaxies belonging to the above defined ‘*Flat*’ and ‘*Merger*’ classes of luminosity profiles (see Section 2). In the left panels the solid lines represent the ‘*local*’ Kormendy relation in the V_{606} band. It was obtained from the $\langle \mu_e \rangle - r_e$ relation, derived in the Gunn- r band using the data of Jørgensen et al.(1995), by shifting the zero point according to the transformations given by Jørgensen (1994) and Holtzman et al. (1995). The size of the symbols in the left panels decreases with increasing redshift.

In the upper panel we plot the $\langle \mu_e \rangle$ as derived from the deconvolved profiles. In the lower panel we have applied to $\langle \mu_e \rangle$ the standard correction due to the cosmological dimming, $(1+z)^4$. The panels on the right show the residuals with respect to the local *KR*, as a function of

the redshift.

A lack of spectroscopic redshifts in the range $1 \lesssim z \lesssim 2$ is apparent. We also note the tendency of high redshift galaxies ($z \gtrsim 2$) to be intrinsically smaller with respect to galaxies with $z \lesssim 1$. By applying only the $(1+z)^4$ cosmological dimming correction (lower-left panel in Figure 4) we obtained two distinct sequences for high redshift and low redshift galaxies in the $\langle \mu_e \rangle - r_e$ plane. Both sequences are roughly parallel to the local KR , with a small negative shift in the zero point at low-redshift and a large positive shift at high redshift. In the following section we investigate how such shifts can be explained in terms of evolutionary effects.

4.1. The K+E-correction: colors and models

In order to derive more realistic corrections for the observed surface brightness of the *HDF* ellipticals, we need to model their evolution.

We have used the chemical-spectrophotometric models of population synthesis produced by Bressan et al. (1994, *BCF*) and Tantalo et al. (1996, *TCBF*) to estimate the K+evolutionary-corrections (K+E) for elliptical galaxies of different mass, age and evolutionary history.

The *BCF* models belong to the ‘*closed-box*’ class. The history of star formation consists of an initial period of activity followed by quiescence after the onset of the galactic winds, whose duration depends on the galactic mass, being longer in the high mass galaxies and shorter in the low-mass ones. In the following we will refer to a *passive* model, characterized by an early onset of the galactic winds, and an *active* model, in which star formation continues until late epochs. They are both described in detail in BCF.

TCBF model the spectrophotometric evolution of elliptical galaxies as governed by the infall scheme. They simulate the collapse of a galaxy made of two components, the luminous and dark material. While the mass of the dark component is supposed to remain constant with time, the mass of the luminous component is assumed to grow at a suitable rate. The effects of galactic winds powered by supernova explosions and stellar winds from massive, early-type stars are taken

into account. We will refer to this scenario as the *infall* models.

In order to verify which subset of the models is able to best describe the elliptical galaxies observed in *HDF* we have compared the observed colors as a function of the (spectroscopic) redshifts with the model predictions. We have obtained from the models the spectral energy distributions (*SED*) of a galaxy at a given redshift as a function of the type of model, galaxy mass, redshift of formation and cosmological parameter q_o . A subset of the results is shown in Fig. 5, where the best-fitting models for each one of the three evolutionary schemes are compared with the observations. It can be seen that a model of a $3 \cdot 10^{12} M_\odot$ galaxy with an ‘*active*’ star formation until the age of 11 Gyr with a redshift of formation $z_{form} = 5$ in a universe with $q_o = 0.5$ provides a reasonable ‘global’ description of the observed colors. At low redshift, two different sequences of galaxy colors are apparent in the $U - V$ (and in minor degree in the $B - V$). To fully reproduce them, different models or at least different choices of the parameters (e.g. z_{form}) characterizing a given model would be required. We have not attempted to do it in this simplified approach, since we were interested only in the K+E corrections to be applied to the V_{606} band. It is remarkable that all the galaxies belonging to the bluer population show ‘*abnormal*’ luminosity profiles (types 2 and 3 in Tab. 1, squares in Fig. 5).

In general, smaller values of q_o can be compensated (although not exactly) by a corresponding variation of the redshift of formation, in the sense that for lower values of q_o the galaxy colors at a given redshift tend to be reproduced by models with smaller z_{form} .

The *passive* closed-box model and the models governed by the infall are clearly not able to reproduce the color- z diagrams with a unique redshift of formation.

In the present analysis we have not included the effects of dust which, due to the shape of the extinction law, are potentially more and more important at increasing redshifts. In general to reproduce the observed colors in presence of dust a smaller redshift of formation is required. On the other hand, a larger unabsorbed luminosity has to be assumed and the two effects tend, at least qualitatively, to compensate in the estimation of the $K + E$ correction.

If we apply to the observed $\langle \mu_e \rangle$ the K+E-corrections predicted by the *BCF active* model with $q_o = 0.5$ and $z_{form} = 5$, we find the K-relation shown in Figure 6, which for all redshifts is in fairly good agreement with the local relation. Again, we note that all galaxies with very high redshift (small symbols) are confined in the high surface brightness, small size region of the plane.

4.2. Extending the sample: photometric redshifts

In order to improve the statistics, we have estimated the photometric redshifts of the 25 galaxies in the present sample having sufficient and reliable infrared information (i.e. detection in J , H and K , no deblending problems), besides the optical magnitudes.

The photometric redshift technique has been successfully used by various authors in the *HDF* (e.g. Lanzetta et al. 1996) and in other fields (e.g. Connolly et al. 1995, Giallongo et al. 1997).

We have adopted an approach of the type described by Giallongo et al.(1997), computing a large number of galaxy SEDs as a function of the formation redshift, redshift of the observation, internal extinction and evolutionary scenario and comparing them with the observed colors. In the present case only population synthesis models of elliptical galaxies (those provided by *BCF* and *TCBF*) were used. In this way, the morphological information allowed us to restrict the range of the practicable models, avoiding possible degeneracies. In practice, the technique exploits at low redshift the position of the 400 nm break and at high-redshift the drop in the galaxy flux due to the combination of the IGM plus intrinsic Lyman absorption.

A complete discussion of the ‘*photometric redshift*’ procedure is beyond the scope of this paper and will be given elsewhere (Arnouts et al. in preparation). In Fig. 7 we show the comparison between the spectroscopic and photometric redshifts for the 23 galaxies in our sample having both redshift determinations.

Fig. 8 shows the KR we obtain for the whole sample, including the galaxies with photometric redshifts (open symbols with error bars) and those with spectroscopic redshifts (same symbols as in Fig. 6). Again the triangles represent the galaxies with ‘*abnormal*’ luminosity profiles and the

size of the symbols is decreasing with increasing redshift.

The KR for galaxies with photometric redshift estimates shows a larger dispersion with respect to the spectroscopic sample. However, the general trend is preserved and the 2-D Kolmogorov–Smirnov test (Fasano and Franceschini 1987) cannot reject the null hypothesis that the two distributions are drawn from the same parent population with at more than 82% significance level. In any case different kinds of biases are expected to affect the two samples (e.g. objects with spectroscopic identifications tend to have on average brighter magnitudes and higher central surface brightnesses).

5. Discussion and Conclusions

Figure 8 indicates that, if we apply to the observed $\langle \mu_e \rangle$ the K+E-correction deduced from the galaxy models that properly reproduce the observed colors, the early-type galaxies in the *HDF* follow the local Kormendy relation at any redshift.

The Kormendy relation is well known to be a projection of the Fundamental Plane (FP) which roughly represents the virial equilibrium condition of galaxies, except for a small tilt interpreted in terms of an $M/L \propto M^\alpha$ relation (Faber et al. 1987).

Steidel et al. (1996) measured the line widths of compact spheroidal galaxies at $z > 3$ which, if interpreted as due to gravitational motions, imply velocity dispersions of the order $180 \leq \sigma \leq 320$ km s⁻¹. Such values would place the high- z galaxies of the present sample in the typical region of the Fundamental Plane corresponding to early-type galaxies, showing that, even at $z > 2$, these galaxies obey an average, rest-frame, virial equilibrium condition. It is worth noticing that, even assuming a significant non-virial contribution (e.g. stellar winds, SN ejecta) to the observed line widths, the high- z galaxies would still be in the FP region occupied by intermediate-luminosity elliptical galaxies.

Fig. 8 also shows that, high redshift galaxies are confined in the high surface brightness, small effective radius region of the KR plane. Is this tendency due to the fact that large ellipticals have

a faint surface brightness (according to the Kormendy relation itself) ? Large galaxies at high redshifts could lie below the detection limit. We have checked at various redshifts what regions of the Kormendy plane result ‘*forbidden*’ because of the adopted selection criteria. The regions of Fig. 8 below the dotted, dashed and dot–dashed lines represent, in the framework of the *BCF* active models, the forbidden regions of the Kormendy plane for $z = 1, 2$ and 3 , respectively. We conclude that, had there been large galaxies at high redshifts in the *HDF*, they would have been included in our sample.

However, it remains to be demonstrated that the comoving volume density of large high redshift galaxies is high enough to ensure the presence of a significant number of them in our sample. We have estimated in two different ways the expected fraction, F_L , of $z > 1.5$ galaxies with $\log(r_e^{kpc}) > 0.2$.

1. Assuming that the r_e distribution at $z > 1.5$ is the same observed in the Coma cluster (Lobo et al. 1997) for galaxies in the same absolute magnitude range, we found $F_L = (166/299) \simeq 1/2$. To carry out this computation, we applied the $K + E$ corrections of the BCF active models and transformed the isophotal radii (at $\mu_V = 26.5$) measured by Lobo et al. (1997) into effective radii by using the average growth curve of the elliptical galaxies given by de Vaucouleurs et al. (1976). It should be noted that the sample by Lobo et al. (1997) contains all morphological types and the distribution of r_e , although ellipticals are preponderant in Coma, may be affected by the presence of spirals and irregulars.
2. In an alternative approach we assumed that the galaxies at $z > 1.5$ - again with the $K + E$ correction derived by the BCF active models - follow a Schechter-type luminosity function with an $M_B^* = -21.7$ (Ferguson and Sandage 1991). Assuming that the luminosity profiles follow the de Vaucouleurs law, we can use the local Kormendy relation to transform the distribution of absolute luminosities in a distribution of effective radii, according to the formula: $-2.5 \log r_e = (M_{606} + 20.15)$. In this way we found again $F_L \simeq 1/2$.

Although both estimates are based on rough approximations, they turn out to be robust

with respect to changes of the models adopted for the $K + E$ corrections, the luminosity function parameterization and possible contaminations of the diameter and/or luminosity distribution functions. Therefore we conclude that at $z > 1.5$ a scarcity of large early-type galaxies is observed with respect to the local universe.

What type of selection effects may affect the present sample ?

The choice of the V_{606} as the selection band implies a bias at redshifts $z > 1$ in favor of star-forming galaxies. So it is not surprising that at $z > 2$ the observed colors of the galaxies favor the ‘active’ models. Galaxies following a ‘passive’ or ‘infall’ evolution, if formed at sufficiently large redshifts, quickly become too faint to be observable. So, a population of redder galaxies which had their major burst of star formation few e-folding times before the epoch of the observations would be undetected in the HDF.

Is also our perspective of the morphology of galaxies possibly biased by the fact that at $z > 1$ the V_{606} band is sampling the ultraviolet part of the SED ? The UV morphological properties of nearby elliptical galaxies are known to match those in the optical (see O’Connell 1997 and references therein), but for these systems the UV light is coming from evolved stars (the UV upturn). On the contrary, in interpreting the morphology of star-forming galaxies at substantial redshifts, one must bear in mind that moving to UV wavelengths the light becomes proportional to the star-formation rates and not to the global stellar content of a galaxy. In principle the UV profiles could depart from an $r^{1/4}$ law, reflecting the comparatively more chaotic morphology of the star formation rather than the structural morphology of a galaxy.

On the other hand, the SED of unreddened star-forming galaxies at high- z is essentially flat from $\sim 1200\text{\AA}$ to $\sim 4000\text{\AA}$ rest frame, and to even longer wavelengths if the galaxy is young. At the wavelength of the V_{606} WFPC2 filter, this corresponds to observing galaxies in the redshift interval $0.5 < z < 4$, with minimal K -corrections (see Giavalisco et al. 1996). This means that it is not unfair to compare the UV morphologies of the $z > 2$ galaxies with those of the $z \sim 1$ systems. In order to check quantitatively this conclusion we produced, in the four WFPC2 available bands, the ‘*aperture*’ luminosity profiles of the galaxies with spectroscopic redshifts

$z < 1.5$ and $U_{300} - V_{606} < 2$. The SED of these galaxies ($3 - 696 - 0$, $2 - 264 - 2$, $4 - 565 - 0$ and $4 - 727 - 0$), obtained from the optical and infrared magnitudes in Table 1, clearly indicates star-forming activity. Nevertheless, the Fig. 9 shows that the light distribution in these objects is ostensibly the same in a wide range of rest-frame wavelengths (from the V band to the far- UV).

6. acknowledgments

We thank A. Bressan, S. Charlot, C. Chiosi and E.V. Held for enlightening discussions and an anonymous referee for stimulating comments. This work was partially supported by the ASI contracts 94-RS-107 and 95-RS-38, by the TMR program of the European Community and by the Formation and Evolution of Galaxies network set up by the European Commission under contract ERB FMRX-CT96-086 of its TMR program.

REFERENCES

- Bender R., Burstein, D., Faber, S. M. 1992, ApJ, 399, 462
- Bender, R., Saglia, R.P, Ziegler, B., Belloni, P., Greggio, L., Hopp, U., Bruzual, G. 1997, astro-ph/9708237
- Bendinelli O., 1991, ApJ, 366, 599
- Bertin E., Arnouts S., 1996, A&AS, 117, 393
- Bressan A., Chiosi, C., Fagotto, F., 1994, ApJS **94**, 63
- Connolly A.J., Csabai I., Szalay A.S., Koo D.C., Kron R.G., Munn J.A., 1995, AJ 110, 2655
- Couch W.J., 1996,
<http://ecf.hq.eso.org/hdf/catalogs/>
- Dickinson M., 1995, Proc. of the workshop: *‘Fresh views of Elliptical Galaxies*, PASP Conference Series, 86, p. 283
- Dickinson M., 1997a, Proc. Third ESO-VLT Workshop: *Galaxy Scaling Relations: Origins, Evolution and Applications*, p.215.
- Dickinson M., 1997b,
<http://www.stsci.edu/ftp/observer/hdf/clearinghouse/clearinghouse.html>
- Djorgovski, S., Davis, M. 1987, ApJ, 313, 59
- Djorgovski, S., Santiago, B.X., 1993, Proc. of the workshop: *Structure, Dynamics and chemical evolution of Elliptical Galaxies*, ESO Proceedings, 45, 59
- Dressler, A., Lynden-Bell, D., Burstein, D., Davies, R. L., Faber, S. M., Terlevich, R. J., Wegner, G. 1987a, ApJ, 313, 42

- Dressler, A., Faber, S. M., Burstein, D., Davies, R. L., Lynden-Bell, D., Terlevich, R. J., Wegner, G. 1987b, ApJ, 313, L37
- Dressler, A., Oemler, A. Jr., Butcher, H., Gunn, J. E. 1994a, ApJ, 430, 107
- Dressler, A., Oemler, A. Jr., Sparks, W. B., Lucas, R. A. 1994b, ApJ, 435, L23
- Ellis, R.S., 1997, Proc. of the STScI May Symp.: ‘*The Hubble Deep Field*, astro-ph/9708075
- Faber S. M., Dressler A., Davies R. L., Burstein D., Lynden-Bell D., Terlevich R. J., Wegner G. 1987, in *Nearly Normal Galaxies*, ed. S. M. Faber, Springer-Verlag, New York, p. 175
- Faber S. M., Wegner G., Burstein D., Davies R. L., Dressler A., Lynden-Bell D., Terlevich R. J. 1989, ApJS, 69, 763
- Fasano, G., Franceschini, A. 1987, MNRAS, 225, 155
- Fasano G., Filippi M., Bertola F., 1996, Proc. Second Stromlo Symposium: *The Nature of Elliptical Galaxies*, PASP Conference Series 116, 557.
- Fasano G., Filippi M., 1998, A&A Suppl., in press (FF98)
- Fasano G., Bettoni D., Kjaergaard P., Moles M., 1997, Proc. Third ESO-VLT Workshop: *Galaxy Scaling Relations: Origins, Evolution and Applications*, p.344.
- Franx M., Illingworth G., 1990, ApJ 359, L41
- Ferguson H.C., Sandage A.R., 1991, AJ 101, 765
- Giallongo E., Charlot S., Cristiani S., D’Odorico S., Fontana A., 1997, in ‘*The Early Universe with VLT*’ (Springer), J. Bergeron Ed., pg. 209
- Giallisco M., Steidel C.C., Macchetto F.D., 1996, ApJ 470, 189
- Jørgensen I., 1994, PASP 106, 967

- Jørgensen I., Franx M., Kjaergaard P., 1995, MNRAS 273, 1097
- Jørgensen I., Hjortl, J., 1997, Proc. Third ESO-VLT Workshop: *Galaxy Scaling Relations: Origins, Evolution and Applications*, p.175.
- Holtzman J.A., Burrows C.J., Casertano S., Hester J.J., Trauger J.T., Watson A.M., Worthey G., 1995, PASP 107, 1065
- Kauffmann G., Charlot S., 1997, submitted to MNRAS, astro-ph/9704148
- Kelson, D.D., van Dokkum, P.G., Franx, M., Illingworth, G.D., Fabricant, D. 1997, ApJ, 478, L13
- Kjaergaard, P., Jørgensen, I., Moles, M. 1993, ApJ, 418, 617
- Kormendy J., 1985, ApJ, 295, 73
- Lanzetta K.M., Yahil A., Fernandez-Soto A., 1996, Nature, 381, 759
- Lilly, S. J., Tresse, L., Hammer, F., Crampton, D., Le Fèvre, O. 1995, ApJ, 455, 108
- Lynden-Bell, D., Faber, S. M., Burstein, D., Davies, R. L., Dressler, A., Terlevich, R. J., Wegner, G. 1988, ApJ, 326, 19
- Lobo C., Biviano A., Durret F., Gerbal D., Le Fevre O., Mazure A., Slezak E., 1997, A&AS 122, 409
- Moore, B., Matz, N., Lake, G., Dressler, A., Oemler, A. Jr. 1996, Nature, 379, 613
- O’Connell R.W., 1997, Proc. of the meeting: ‘*The Ultraviolet Universe at Low and High Redshift*’, in press, astro-ph/9706265
- Pahre M.A., Djorgovski S.G. 1996, Proc. of the ‘*Second Stromlo Symposium*’, PASP Conference Series, 116, p. 154
- Pahre M.A., Djorgovski S.G., de Carvalho R.R. 1996, ApJ 456, L79

- Peletier R.F., Davies R.L., Davis L.E., Illingworth G.D., Cawson M., 1990, AJ 100, 1091
- Statler T., 1996, private communication
- Steidel C. C., Giavalisco M., Pettini M., Dickinson M., Adelberger K., 1996, ApJ, 462, 17
- Tantalo R., Chiosi C., Bressan A., Fagotto F., 1996, A&A 311, 361
- Tully, R.B., Fisher, J.R.T., 1977, A&A 54, 661
- van den Bergh S., Abraham R.G., Ellis R.S., Tanvir N.R., Santiago B.X., Glazebrook K.G., 1996, AJ 112, 359
- van Dokkum, P.G., Franx, M. 1996, MNRAS, 281, 958
- de Vaucouleurs G., de Vaucouleurs A., Corwin H.G., 1976, Second Reference Catalogue of Bright Galaxies, University of Texas press, Austin and London.
- Williams R. E., Blacker B., Dickinson M., Dixon W., Ferguson H. C., Fruchter A. S., Giavalisco M., Gilliland R. L., Heyer I., Katsanis R., Levay Z., Lucas R. A., Mcelroy D. B., Petro L., Postman M., 1996, AJ, 112, 1335




Fig. 1.— Difference between the magnitudes obtained with detailed surface photometry and the SExtractor aperture magnitudes vs. the observed average surface brightness (right panel) and the effective radius (left panel).




Fig. 2.— Examples of luminosity profile deconvolution of large (upper panels) and small (lower panels) HDF ellipticals. The solid lines closely following the observed profiles (small dots) represent the re-convolutions of the deconvolved profiles (solid lines with brighter luminosity peaks). The dashed lines in each panel illustrate the multigaussian decomposition of the deconvolved profiles.




Fig. 3.— Comparison between the effective radii before and after deconvolution for the HDF ellipticals in our sample (open circles) and for a 10×10 grid of toy models of different sizes and luminosities. Each toy model was obtained by adding the proper noise to a synthetic elliptical galaxy ($r^{1/4}$ luminosity profile) convolved with the HDF *PSF*. For each value of the ‘true’ effective radius in the grid, 10 toy models with different luminosities were produced and each toy model was processed in the same way as the HDF ellipticals (detailed surface photometry and deconvolution). The solid line connects the average values of r_e measured before the deconvolution as a function of the ‘true’ r_e . The filled circles represent the corresponding average values of r_e obtained after the deconvolution. Error bars are reported in both cases.




Fig. 4.— The μ_e - r_e plane for HDF ellipticals with spectroscopic redshift (left panels). In the upper panel we plot the $\langle \mu_e \rangle$ as derived from the deconvolved profiles. The average error bars in both coordinates are reported in the upper-right part of the plot. In the lower panel we have applied to $\langle \mu_e \rangle$ the standard correction for the cosmological dimming, $(1+z)^4$. The solid line represents the local KR in the V_{606} band. The triangles represent the galaxies belonging to the classes 2 or 3 of luminosity profiles. The size of the symbols decreases with increasing redshift. The panels on the right show the residuals with respect to the local KR as a function of the redshift.




Fig. 5.— Colors of HDF ellipticals with spectroscopic redshift in our sample as a function of the redshift. The symbols are the same as in fig. 4. The solid, dotted and dashed lines represent the BCF-active, BCF-passive and TCBF evolutionary models respectively. Upper and lower limits are represented by one-sided bars.




Fig. 6.— The μ_e - r_e relation for HDF ellipticals with spectroscopic redshift in our sample (left panel). Symbols are the same as in fig. 4. In this case we have applied to the observed $\langle \mu_e \rangle$ the K+E-corrections predicted by the *BCF active* model with $q_o = 0.5$ and $z_{form} = 5$. The panel on the right shows the residuals with respect to the local *KR*, as a function of the redshift.




Fig. 7.— Comparison between the spectroscopic and photometric redshifts.




Fig. 8.— The Kormendy relation for HDF ellipticals in our sample. The filled and empty symbols refer to galaxies with spectroscopic and photometric redshift, respectively. The symbols and the symbol sizes are the same as in fig. 4. The error bars of galaxies with photometric redshift are due to the uncertainty in the redshift estimate, as well as to the average errors of r_e and $\langle \mu_e \rangle$. In the framework of the *BCF* active models, the regions below the dotted, dashed and dot-dashed lines represent the regions of the Kormendy plane which are ‘forbidden’ because of the adopted selection criteria, for $z = 1, 2$ and 3 , respectively.




Fig. 9.— Comparison among the aperture luminosity profiles in the four WFPC bands for the four star-forming galaxies in our sample with spectroscopic redshift $0.4 < z < 1.25$. The surface brightness is expressed in arbitrary units and the central values of μ in the four bands are shifted each other by $0^m.5$.

TABLE 1. Ellipticals sample in HDF

ID-Focas	R.A.(2000)		Dec.(2000)		Type	r_e	$<\mu_e>$	U_{AB}	B_{AB}	V_{AB}	I_{AB}	J	H	K	Note	z	Δz
	'	"	'	"	(1)	"		(2)	(2)	(3)	(2)	(4)	(4)	(4)	(5)	(6)	
4-942-0	36	39.43	12	11.76	1	0.44	24.97	26.13	25.58	24.55	23.10	21.32	20.22	19.54	2	(0.76)	0.08
4-926-2	36	39.56	12	13.83	3	0.13	22.93	<27.99	26.32	25.16	25.36	21.48	20.32	19.53	3	(2.09)	0.25
4-928-1	36	40.01	12	7.37	1	0.32	23.10	26.08	24.60	23.41	22.06	20.12	19.48	18.62	0	1.01 ^a	-
4-878-11	36	40.96	12	5.31	2	0.16	21.72	25.68	23.89	23.46	23.17	21.32	20.68	19.82	2	(1.92)	0.02
4-858-13	36	41.25	12	3.07	1	0.08	21.81	<27.99	25.75	24.98	24.72	22.23	21.95	20.66	6	3.23 ^b	-
4-767-0	36	41.49	12	14.98	2	0.13	22.76	<27.99	26.83	24.96	23.29	20.95	20.11	19.16	1	(0.81)	0.12
4-639-1	36	41.71	12	38.75	1	0.10	22.12	<27.99	24.91	24.92	24.75	<23.43	<22.36	<21.93	1	2.59 ^b	-
4-937-0	36	42.28	11	26.18	1	0.12	22.92	26.73	26.49	25.30	24.94	<23.43	<22.36	<21.93	1	(0.56)	0.13
1-95-0	36	42.38	13	19.36	1	0.07	21.23	25.41	25.14	24.83	23.91	22.76	22.05	21.43	0	(0.87)	0.08
4-493-0	36	43.16	12	42.20	1	0.41	23.30	27.34	25.14	23.06	21.40	19.60	18.74	17.90	0	0.85 ^a	-
4-727-0	36	43.41	11	51.57	2	0.16	21.51	23.59	23.31	23.23	22.85	21.47	20.65	19.87	2	1.24 ^d	-
4-565-0	36	43.63	12	18.25	3	0.26	22.53	24.07	23.74	23.26	22.43	21.04	20.43	19.93	0	0.75 ^a	-
4-744-0	36	43.80	11	42.88	1	0.57	23.09	26.20	24.18	22.13	20.53	18.91	18.06	17.28	2	0.76 ^a	-
2-82-1	36	44.07	14	9.92	2	0.13	22.42	26.41	24.74	24.72	24.67	<23.43	<22.36	<21.93	1	2.27 ^c	-
4-752-1	36	44.38	11	33.20	1	0.84	24.59	27.26	25.01	22.77	20.88	18.66	17.73	16.82	2	1.01 ^a	-
4-555-2	36	45.33	11	54.52	1	0.27	23.97	27.55	26.36	24.59	22.99	20.20	19.42	18.55	6	(1.22)	0.12
1-35-0	36	45.61	13	8.92	1	0.14	22.14	25.39	24.92	24.15	23.52	22.21	21.52	20.90	0	(0.64)	0.08
4-254-0	36	46.13	12	46.50	1	0.47	24.01	26.34	24.83	23.44	21.90	20.30	19.55	18.86	0	(0.81)	0.10
1-47-0	36	46.16	13	13.89	2	0.09	22.98	25.76	25.39	24.97	24.25	23.43	<22.36	<21.93	1	(0.69)	0.15
4-322-2	36	46.21	12	28.43	2	0.11	22.71	26.73	25.57	25.39	24.97	22.72	21.32	21.45	0	(1.95)	0.38
2-251-0	36	46.34	14	4.62	1	0.42	23.09	24.95	23.55	22.76	21.53	19.14	18.41	17.57	0	0.96 ^a	-
4-471-0	36	46.51	11	51.32	1	0.17	21.27	27.01	24.57	22.91	21.86	20.27	19.53	18.86	0	0.51 ^a	-
4-289-0	36	46.95	12	26.08	1	0.09	22.57	<27.99	25.36	25.53	25.17	<23.43	<22.36	<21.93	1	(2.97)	0.11
2-201-0	36	47.18	13	41.82	1	0.08	20.92	25.18	24.41	24.25	23.75	21.76	21.37	20.66	2	(1.32)	0.08
2-121-111	36	48.08	13	9.02	1	0.52	22.07	25.91	23.06	21.28	20.24	18.76	17.93	17.24	0	0.47 ^a	-
2-412-11	36	48.11	14	14.42	1	0.12	22.74	25.85	25.61	25.22	24.51	22.46	21.94	20.86	2	(1.27)	0.10
2-449-1	36	48.34	14	16.63	2	0.17	22.26	25.63	23.90	23.91	23.39	21.35	20.58	19.91	2	2.01 ^d	-
2-173-0	36	48.47	13	16.62	2	0.64	24.76	27.93	24.81	23.54	22.69	21.28	20.03	19.86	2	(0.54)	0.10
2-180-0	36	49.05	13	9.64	2	0.15	22.57	25.21	24.61	24.45	23.77	23.10	<22.36	<21.93	3	(0.86)	0.17
2-264-2	36	49.38	13	11.22	1	0.14	20.56	23.77	23.22	22.60	22.03	20.57	19.70	19.23	2	0.48 ^a	-
2-456-1111	36	49.44	13	46.88	1	0.46	19.24	22.72	19.77	18.73	18.09	16.63	15.97	15.59	0	0.09 ^a	-
3-229-0	36	49.48	12	48.73	2	0.19	23.30	26.16	25.53	24.76	23.79	22.63	22.15	21.26	0	(0.73)	0.06
3-143-0	36	49.64	12	57.43	3	0.29	22.15	24.74	23.73	22.65	21.92	20.25	19.31	18.61	0	0.48 ^a	-
3-243-0	36	49.81	12	48.79	1	0.14	23.31	<27.96	26.29	25.32	24.94	<23.43	<22.36	<21.93	1	3.23 ^c	-
3-321-1	36	50.27	12	45.75	1	0.36	22.85	26.95	24.82	22.85	21.36	19.43	18.64	17.88	0	0.68 ^a	-
3-586-0	36	52.10	12	26.31	1	0.29	25.08	<27.96	27.35	25.57	23.79	21.28	20.20	19.30	1	(1.06)	0.10

TABLE 1. (continued)

ID-Focas	R.A.(2000)		Dec.(2000)		Type	r_e	$\langle \mu_e \rangle$	U_{AB}	B_{AB}	V_{AB}	I_{AB}	J	H	K	Note	z	Δz
	'	"	'	"	(1)	"		(2)	(2)	(3)	(2)	(4)	(4)	(4)	(5)	(6)	
3-696-0	36	52.69	12	19.72	2	0.15	21.64	24.76	24.37	23.57	23.11	21.72	22.03	20.72	0	0.40 ^a	-
2-637-0	36	52.76	13	39.08	1	0.10	22.26	<28.00	25.84	24.98	24.78	<23.43	<22.36	<21.93	1	3.37 ^c	-
2-591-2	36	53.18	13	22.75	2	0.13	22.54	<28.00	25.21	24.85	24.73	<23.43	<22.36	<21.93	1	2.49 ^d	-
2-643-0	36	53.42	13	29.52	3	0.19	23.40	<28.00	25.70	24.83	24.55	<23.43	<22.36	<21.93	1	2.99 ^c	-
3-118-1	36	54.73	13	14.73	1	0.09	21.26	26.10	24.58	24.39	24.27	21.86	21.01	20.85	2	(2.25)	0.12
2-898-0	36	54.78	13	50.74	3	0.23	23.97	25.75	25.36	24.94	24.02	22.86	<22.36	<21.93	1	(0.86)	0.20
3-266-0	36	55.16	13	3.60	1	0.16	23.19	<27.96	27.08	24.99	23.30	21.25	20.46	19.65	1	(0.73)	0.10
3-180-1	36	55.46	13	11.19	1	0.47	24.29	27.94	25.53	23.74	21.99	19.67	18.83	17.92	0	(0.99)	0.10
3-904-0	36	55.95	12	10.72	1	0.10	22.08	25.82	25.08	24.91	24.39	22.45	<22.36	21.51	1	(1.27)	0.12
3-815-1	36	56.65	12	20.12	1	0.36	24.06	26.88	26.04	24.09	22.39	20.26	19.42	18.58	0	(0.81)	0.12
3-355-0	36	56.92	13	1.56	1	0.33	23.78	24.92	24.29	23.96	22.75	20.00	19.33	18.36	2	(1.23)	0.10
3-790-1	37	0.56	12	34.60	1	0.35	22.30	26.21	24.08	22.40	21.26	19.74	18.92	18.21	0	0.56 ^a	-

Notes to Table 1.

Notes :

⁽¹⁾classification : 1= normal; 2= intermediate; 3 = Merger⁽²⁾computed by using adaptive aperture colours in AB system⁽³⁾Luminosity profile magnitude in AB system⁽⁴⁾total magnitude performed with SExtractor in standard zero-point system⁽⁵⁾Classification by eyes : 0= isolated galaxie; 1 = at least one upper limit ; 2 = neighbours can be affected the magnitudes ;4 = Merger in IR (J,H,K) ; final flag is FLAG = 0 + 2⁰ + 2¹ + 2²⁽⁶⁾spectroscopic redshifts come from : (a) Cohen et al. (1996); (b) Steidel et al. (1996) (c) Lowenthal et al. (1997); (d) Dickinson et al. (1997).The photometric redshifts (in parentheses) are derived from the optical *and* Infra Red bands